

---

Dilluns 13 de gener del 2014, 15:00h

Aula Petita (CRM).

---

## Geometric-type Sobolev inequalities and applications to the regularity of extremal solutions

MANEL SANCHON

(Universitat Autònoma de Barcelona)

**ABSTRACT:** We will consider the reaction diffusion equation  $-\Delta u = \lambda f(u)$  with zero Dirichlet boundary condition, where  $\Omega \subset \mathbb{R}^n$  is a smooth bounded domain,  $\lambda$  is a positive parameter, and the reaction term  $f \in C^1(\mathbb{R})$  is an increasing function satisfying  $f(0) > 0$  and  $f(t)/t \rightarrow +\infty$  as  $t \rightarrow +\infty$ . Under these assumptions it is well known that there exists an extremal parameter  $\lambda^* \in (0, +\infty)$  such that this problem admits a classical (minimal) solution  $u_\lambda$  for  $\lambda \in (0, \lambda^*)$  and admits no weak solution for  $\lambda > \lambda^*$ . The increasing limit  $u^* := \lim_{\lambda \uparrow \lambda^*} u_\lambda$  is known to be a weak solution for  $\lambda = \lambda^*$  and it is called the extremal solution. The regularity of  $u^*$  has been an active topic of research in the last decade. However for general reaction terms  $f$  and general domains  $\Omega$  the known results are far to be optimal.

In this talk I will introduce some of the results available in the literature concerning the regularity of the extremal solution and I will concentrate my attention in some  $L^q$  and  $W^{1,q}$  estimates obtained recently in [1] and [2]. In order to obtain such estimates we will prove some geometric Sobolev inequalities involving the mean curvature of the function appearing in the inequalities.

## References

- [1] X. Cabré and M. Sanchón, Geometric-type Sobolev inequalities and applications to the regularity of minimizers, *J. Funct. Anal.* **264** (2013), 303-325.
- [2] M. Sanchón,  $W^{1,q}$  estimates for the extremal solution of reaction-diffusion problems, *Nonlinear analysis* **80** (2013), 49,54.